| P | llack | Rutto | Conner | Project | Mino | Operating | Darmit | Application | (Pavision ' | 21 |
|---|--------|-------|--------|----------|--------|-----------|--------|--------------|----------------|------------|
| ப | iiaun. | Dulle | CUDDEI | 1 101001 | IVIIII | ODELALITU | | ADDIIGALIOLI | II VEVISIUII A | <i>_ 1</i> |

APPENDIX N-1: Post Closure Nondegredation Evaluation

Tintina Montana, Inc.

May 2017

MEMORANDUM

DATE: June 10, 2016

TO: Bob Jacko, Tintina Montana

Allen Kirk, Geomin Resources

FROM: Greg Bryce, Project Manager

James Lloyd, P.E.

SUBJECT: Post Closure Nondegradation Evaluation, Black Butte Copper Project

Hydrometric has conducted an evaluation of nondegradation criteria for Post Closure water quality flowing through the mine working of the Upper Copper Zone (UCZ) at the Black Butte Copper Project. The UCZ mine workings are proposed to be developed in four hydrostratigraphic units: Upper Newland Shales (Ynl A), Upper Sulfide Zone (USZ), Upper Copper Zone (UCZ) and the Lower Newland Shales (Ynl B). The nondegradation analysis was conducted from wells completed within these hydro-stratigraphic units in the vicinity of the UCZ as discussed below and listed in Table 1. The nondegradation criteria were evaluated based on the following elements listed below:

- Ambient groundwater conditions are based on wells completed in the Ynl A, USZ, UCZ, and Ynl B. Water quality data from wells MW-1B, MW-2A, MW-2B, MW-3, MW-9, PW-2, PW-3, PW-4, PW-8, PW-9, and PW-10 were combined; statistics were generated for the 25th, mean and 75th percentiles values. In most cases the 75th percentile value was used to represent ambient water quality.
- Only those parameters that have groundwater standards were evaluated. No additional human health advisories were incorporated into the analysis, however this may occur during the MEPA process.
- The development of the nondegradation criteria used the procedures found in ARM17.30.715. Based on the parameter category the criteria were developed as:
 - No increase allowed in the receiving water for carcinogens and toxics with a BCF>300.
 - An increase (in the receiving water) of 15% of the most stringent water quality standard for toxic parameters.
 - An increase (in the receiving water) of 10% of the most stringent water quality standard for harmful parameters.

 When water quality standards were not promulgated, yet a trigger value was available, the ambient water quality and trigger value were used to set the nondegradation criteria.

Compliance with the nondegradation nonsignificant policy is evaluated by comparing the resulting water quality at closure to the nondegradation criteria calculated by summing the preexisting water quality and the nondegradation criteria. The statistic of the groundwater data and nondegradation evaluation is summarized in Table 1.

To meet the non-significance criteria at closure the following water management option are available:

- Request a regulatory mixing zone (standard or source specific) for ambient mixing of groundwater and waters generated in the mine void. Care should be taken to exclude any influences to surface waters prior to achieving the nondegradation criteria.
- Additives could be included in the paste backfill to limit the leaching of constituents (if any).

Table 1. Estimated Groundwater Non-Degradation Criteria for Underground Workings at Closure

| Constituent | | Groundwater Wells (MW-1B, MW-2A, MW-2B, MW-3, MW-9, PW-2, PW-3, PW-4, PW-8, PW-9, and PW-10) | | | Groundwater Human Health Standard | Category | Non Deg Trigger Level | Ambient + Trigger | Applicable Nonsignificance Factor ARM 17.30.715 | Non Deg Threshold | Required Reporting Limit (RRL) | Estimated Non-Degradation Criteria |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------|----------------------------------------------------------------------------------------------------|----------|----------|-----------------------------------------|-------------------|--------------------------|----------------------|-------------------------------------------------|----------------------|--------------------------------------|------------------------------------------|
| ANTIMONY (Sb) | Constituent | 25%ile | 75%ile | Average | | | | | AKWI 17.30.715 | | | |
| ARSENIC (AS) | ` / | | | | | | | | | | | |
| BABRUM (Ba) | ANTIMONY (Sb) | < 0.0005 | | | 0.006 | Toxic | | | | | | 0.002 |
| | ARSENIC (As) | | | | 0.010 | Carcinogen | | | | NAI | | 0.064 |
| CADMUM (Cd) | BARIUM (Ba) | 0.013 | 0.043 | 0.035 | 1.0 | Toxic | 0.002 | 0.0448 | 0.15 | 0.1500 | 0.005 | 0.193 |
| CHROMIUM (C) | BERYLLIUM (Be) | < 0.0008 | 0.0010 | 0.0009 | 0.004 | Carcinogen | NAI | NAI | NAI | NAI | 0.001 | 0.0010 |
| COBALT (CO) | CADMIUM (Cd) | < 0.00003 | 0.00007 | 0.00004 | 0.005 | Toxic | 0.0001 | 0.0002 | 0.15 | 0.0008 | 0.00008 | 0.00082 |
| COPPER (Cu) | CHROMIUM (Cr) | < 0.001 | 0.010 | 0.006 | 0.1 | Toxic | 0.001 | 0.0110 | 0.15 | 0.0150 | 0.001 | 0.025 |
| FLUGRIDE (F) | COBALT (Co) | < 0.01 | 0.01 | 0.01 | | | | | | | 0.01 | |
| RON (Fe) | | | | | | Toxic | 0.0005 | | | | | 0.197 |
| IEAD (Pb) | FLUORIDE (F) | 0.30 | 0.60 | 0.43 | 4.0 | Toxic | 0.005 | 0.6050 | 0.15 | 0.6000 | 0.001 | 1.20 |
| MANGANESE (Mn) 0.012 0.085 0.052 0.005 MERCURY (Hg) <0.000005 0.000010 0.000007 0.002 Toxic w/BCF>300 NAI < | IRON (Fe) | 0.068 | 2.140 | 5.513 | | Harmful | | | | | 0.05 | |
| MERCURY (Hg) | LEAD (Pb) | < 0.0003 | 0.0005 | 0.0015 | 0.015 | Toxic | 0.0001 | 0.0006 | 0.15 | 0.00225 | 0.00050 | 0.0028 |
| MOLYBDENUM (Mo) | MANGANESE (Mn) | 0.012 | 0.085 | 0.052 | | | | | | | 0.005 | |
| NICKEL (Ni) 0.001 0.010 0.005 0.1 Toxic 0.0005 0.0105 0.15 0.0150 0.010 | MERCURY (Hg) | < 0.000005 | 0.000010 | 0.000007 | 0.002 | Toxic w/ BCF >300 | NAI | NAI | NAI | NAI | 0.00001 | 0.000010 |
| SELENIUM (Se) 0.0002 0.0010 0.0006 0.05 Toxic 0.0006 0.016 0.15 0.0075 0.0010 | MOLYBDENUM (Mo) | < 0.002 | 0.005 | 0.003 | | | | | | | 0.002 | |
| SILVER (Ag) <0.0005 0.0200 0.0106 0.1 Toxic 0.0002 0.0202 0.15 0.0150 0.001 STRONTIUM (Sr) 0.10 6.48 3.84 4.0 Toxic 0.1 6.5800 0.15 0.6000 0.0002 THALLIUM (TI) 0.0004 0.0039 0.0040 0.002 Toxic 0.0003 0.0042 0.15 0.6000 0.0002 URANIUM (U) 0.0010 0.0080 0.0047 0.03 Carcinogen NAI NAI <td< td=""><td>NICKEL (Ni)</td><td>0.001</td><td>0.010</td><td>0.005</td><td>0.1</td><td>Toxic</td><td>0.0005</td><td>0.0105</td><td>0.15</td><td>0.0150</td><td>0.010</td><td>0.025</td></td<> | NICKEL (Ni) | 0.001 | 0.010 | 0.005 | 0.1 | Toxic | 0.0005 | 0.0105 | 0.15 | 0.0150 | 0.010 | 0.025 |
| STRONTIUM (Sr) 0.10 6.48 3.84 4.0 Toxic 0.1 6.5800 0.15 0.6000 0.0002 THALLIUM (TI) 0.0004 0.0039 0.0040 0.002 Toxic 0.0003 0.0042 0.15 0.0003 0.0002 URANIUM (U) 0.0010 0.0080 0.0047 0.03 Carcinogen NAI | SELENIUM (Se) | 0.0002 | 0.0010 | 0.0006 | 0.05 | Toxic | 0.0006 | 0.0016 | 0.15 | 0.0075 | 0.0010 | 0.009 |
| THALLIUM (TI) | SILVER (Ag) | < 0.0005 | 0.0200 | 0.0106 | 0.1 | Toxic | 0.0002 | 0.0202 | 0.15 | 0.0150 | 0.001 | 0.0350 |
| URANIUM (U) 0.0010 0.0080 0.0047 0.03 Carcinogen NAI NAI NAI NAI 0.008 ZINC (Zn) 0.002 0.01675 0.01573 2.0 Toxic 0.005 0.0218 0.15 0.3000 0.01 NITRATE + NITRITE AS N 0.0100 0.0450 0.0637 10 Toxic 7.5 7.5 0.15 7.5 0.01 NITROGEN (N) TOT 0.5 0.5 0.5 Nutrient 0.01 PHOSPHORUS (P) TOT 0.0 0.0 0.0 Nutrient 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 0.001 <td>STRONTIUM (Sr)</td> <td>0.10</td> <td>6.48</td> <td>3.84</td> <td>4.0</td> <td>Toxic</td> <td>0.1</td> <td>6.5800</td> <td>0.15</td> <td>0.6000</td> <td>0.0002</td> <td>6.48</td> | STRONTIUM (Sr) | 0.10 | 6.48 | 3.84 | 4.0 | Toxic | 0.1 | 6.5800 | 0.15 | 0.6000 | 0.0002 | 6.48 |
| ZINC (Zn) | THALLIUM (TI) | 0.0004 | 0.0039 | 0.0040 | 0.002 | Toxic | 0.0003 | 0.0042 | 0.15 | 0.0003 | 0.0002 | 0.0039 |
| NITRATE + NITRITE AS N 0.0100 0.0450 0.0637 10 Toxic 7.5 7.5 0.15 7.5 0.01 NITROGEN (N) TOT 0.5 0.5 0.5 0.5 Nutrient 0.01 PHOSPHORUS (P) TOT 0.0 0.0 0.0 0.0 Nutrient 0.001 0.001 PH FLD (S.U.) 6.5 7.3 6.9 6.5-8.5 Harmful 0.5 s.u 1 SPECIFIC CONDUCTIVITY (UMHOS/CM) 451.0 751.0 592.7 <1000 Class I GW | URANIUM (U) | 0.0010 | 0.0080 | 0.0047 | 0.03 | Carcinogen | NAI | NAI | NAI | NAI | 0.008 | 0.0080 |
| NITROGEN (N) TOT | ZINC (Zn) | 0.002 | 0.01675 | 0.01573 | 2.0 | Toxic | 0.005 | 0.0218 | 0.15 | 0.3000 | 0.01 | 0.32 |
| PHOSPHORUS (P) TOT 0.0 0.0 0.0 Nutrient 0.001 0.001 PH FLD (S.U.) 6.5 7.3 6.9 6.5-8.5 Harmful 0.5 s.u. 1 1 2 1 250* SMCL < | NITRATE + NITRITE AS N | 0.0100 | 0.0450 | 0.0637 | 10 | Toxic | 7.5 | 7.5 | 0.15 | 7.5 | 0.01 | 7.5 |
| PH FLD (S.U.) 6.5 7.3 6.9 6.5-8.5 Harmful 0.5 s.u 0.5 SPECIFIC CONDUCTIVITY (UMHOS/CM) 451.0 751.0 592.7 <1000 Class I GW 1 SULFATE (SO4) 39 230 146 250* SMCL 1 CHLORIDE (CI) 1 2 1 250* SMCL 1 CALCIUM (Ca) 52 79 63 1 SODIUM (Na) 3 10 6 | NITROGEN (N) TOT | 0.5 | 0.5 | 0.5 | | Nutrient | | | | | 0.01 | |
| SPECIFIC CONDUCTIVITY (UMHOS/CM) 451.0 751.0 592.7 <1000 Class I GW 1 SULFATE (SO4) 39 230 146 250* SMCL 1 CHLORIDE (CI) 1 2 1 250* SMCL 1 CALCIUM (Ca) 52 79 63 1 SODIUM (Na) 3 10 6 1 MAGNESIUM (Mg) 28 49 36 1 TOTAL ALKALINITY AS CaCO3 190.0 220.0 185.8 | PHOSPHORUS (P) TOT | 0.0 | 0.0 | 0.0 | | Nutrient | 0.001 | | | | 0.001 | |
| SULFATE (SO4) 39 230 146 250* SMCL 1 CHLORIDE (CI) 1 2 1 250* SMCL 1 CALCIUM (Ca) 52 79 63 1 SODIUM (Na) 3 10 6 1 MAGNESIUM (Mg) 28 49 36 1 TOTAL ALKALINITY AS CaCO3 190.0 220.0 185.8 10 TOTAL SUSPENDED SOLIDS (TSS) 10.0 10.0 20.0 | PH FLD (S.U.) | 6.5 | 7.3 | 6.9 | 6.5-8.5 | Harmful | 0.5 s.u. | | | | | 6.0 -7.8 |
| CHLORIDE (CI) 1 2 1 250* SMCL 1 CALCIUM (Ca) 52 79 63 1 SODIUM (Na) 3 10 6 1 MAGNESIUM (Mg) 28 49 36 1 TOTAL ALKALINITY AS CaCO3 190.0 220.0 185.8 4 TOTAL SUSPENDED SOLIDS (TSS) 10.0 10.0 20.0 10 | SPECIFIC CONDUCTIVITY (UMHOS/CM) | 451.0 | 751.0 | 592.7 | <1000 | Class I GW | | | | | 1 | <1000 |
| CALCIUM (Ca) 52 79 63 1 SODIUM (Na) 3 10 6 1 MAGNESIUM (Mg) 28 49 36 1 TOTAL ALKALINITY AS CaCO3 190.0 220.0 185.8 4 TOTAL SUSPENDED SOLIDS (TSS) 10.0 10.0 20.0 10 | SULFATE (SO4) | 39 | 230 | 146 | 250* | SMCL | | | | | 1 | 250* |
| SODIUM (Na) 3 10 6 1 MAGNESIUM (Mg) 28 49 36 1 TOTAL ALKALINITY AS CaCO3 190.0 220.0 185.8 4 TOTAL SUSPENDED SOLIDS (TSS) 10.0 10.0 20.0 10 | CHLORIDE (CI) | 1 | 2 | 1 | 250* | SMCL | | | | | 1 | 250* |
| MAGNESIUM (Mg) 28 49 36 1 TOTAL ALKALINITY AS CaCO3 190.0 220.0 185.8 4 TOTAL SUSPENDED SOLIDS (TSS) 10.0 10.0 20.0 10 | CALCIUM (Ca) | 52 | 79 | 63 | | | | | | | 1 | |
| TOTAL ALKALINITY AS CaCO3 190.0 220.0 185.8 4 TOTAL SUSPENDED SOLIDS (TSS) 10.0 10.0 20.0 10 | SODIUM (Na) | | 10 | | | | | | | | 1 | |
| TOTAL SUSPENDED SOLIDS (TSS) 10.0 10.0 20.0 10 | MAGNESIUM (Mg) | 28 | 49 | 36 | | | | | | | 1 | |
| | TOTAL ALKALINITY AS CaCO3 | 190.0 | 220.0 | 185.8 | | | | | | | 4 | |
| TOTALY VIA DEVICE A G. CO.2 | TOTAL SUSPENDED SOLIDS (TSS) | 10.0 | 10.0 | 20.0 | | | | | | | 10 | |
| TOTAL HARDNESS AS CaCO3 244 405 306 | TOTAL HARDNESS AS CaCO3 | 244 | 405 | 306 | | | | | | | | |

NAI = No Allowable Increase (applies to all Carcinogen and Toxics with BCF >300); -- = Not Applicable
Statistics calculated using the value of detection limit when less that detection results. Average value assigned < when 50% or more of samples below detect.

BCF = bio-concentration factor

^{*} Based on EPA Secondary Standard (SMCL)