

**Appendix A: Phosphorus Index Worksheet (Complete for each field and crop)**

Field: B&H1		Crop: Barley/Winter Wheat				Year: 2020		
Field Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0,1,2,4,8)	Weight Factor	Weight Risk
Soil Erosion	NA	<5 tons/as/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	QA> 10 for erodible soils	0	1.5	0.0
Furrow Irrigation Erosion	N/A	Tail water recovery, QS>6 very erodible soils, or QS>10 other soils	QS> for erosion resistant soil	QS> for erodible soils	QA>6 for very erodible soils	0	1.5	0
Sprinkler Irrigation Erosion	All fields 0-3% slope, all sandy fields or field evaluation indicates little or no runoff large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8% large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	1.5	0.0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1 <input type="text"/>	0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	0	0.5	0.0
Commercial P Fertilizer Application Method	None Applied	Placed with Planter or injection deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporate >3 months before crop or surface applied <3 months before crop emerges	Surface applied to pasture or >3 months before crop emerges	1 <input type="text"/>	1.0	1.0
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	1 <input type="text"/>	1.0	1.0

Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges	0	1.0	0.0
Organic P Source Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	0	1.0	0.0
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or application are directly into concentrated surface water flow areas.	0	1.0	0.0
<b>Total Phosphorus Index Value:</b>								<b>2.5</b>

**Appendix A: Phosphorus Index Worksheet (Complete for each field and crop)**

Field: Dorey1								
Crop: Alfalfa Grass Mix								
Year: 2020								
Field Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0,1,2,4,8)	Weight Factor	Weight Risk
Soil Erosion	NA	<5 tons/as/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	QA> 10 for erodible soils	0	1.5	0.0
Furrow Irrigation Erosion	N/A	Tail water recovery, QS>6 very erodible soils, or QS>10 other soils	QS> for erosion resistant soil	QS> for erodible soils	QA>6 for very erodible soils	0	1.5	0
Sprinkler Irrigation Erosion	All fields 0-3% slope, all sandy fields or field evaluation indicates little or no runoff large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8% large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	1.5	0.0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1 <input type="text"/>	0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	0	0.5	0.0
Commercial P Fertilizer Application Method	None Applied	Placed with Planter or injection deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporate >3 months before crop or surface applied <3 months before crop emerges	Surface applied to pasture or >3 months before crop emerges	8 <input type="text"/>	1.0	8.0
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	1 <input type="text"/>	1.0	1.0

Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges	0	1.0	0.0
Organic P Source Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	0	1.0	0.0
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or application are directly into concentrated surface water flow areas.	0	1.0	0.0
<b>Total Phosphorus Index Value:</b>								<b>9.5</b>

### Appendix A: Phosphorus Index Worksheet (Complete for each field and crop)

Field: Dorey2		Crop: Winter Wheat				Year: 2020		
Field Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0,1,2,4,8)	Weight Factor	Weight Risk
Soil Erosion	NA	<5 tons/as/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	QA> 10 for erodible soils	0	1.5	0.0
Furrow Irrigation Erosion	N/A	Tail water recovery, QS>6 very erodible soils, or QS>10 other soils	QS> for erosion resistant soil	QS> for erodible soils	QA>6 for very erodible soils	0	1.5	0
Sprinkler Irrigation Erosion	All fields 0-3% slope, all sandy fields or field evaluation indicates little or no runoff large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8% large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	1.5	0.0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1 <input type="button" value="v"/>	0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	0	0.5	0.0
Commercial P Fertilizer Application Method	None Applied	Placed with Planter or injection deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporate d >3 months before crop or surface applied <3 months before crop emerges	Surface applied to pasture or >3 months before crop emerges	1 <input type="button" value="v"/>	1.0	1.0
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	1 <input type="button" value="v"/>	1.0	1.0

Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges	0	1.0	0.0
Organic P Source Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	0	1.0	0.0
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or application are directly into concentrated surface water flow areas.	0	1.0	0.0
<b>Total Phosphorus Index Value:</b>								<b>2.5</b>

**Appendix A: Phosphorus Index Worksheet (Complete for each field and crop)**

Field: Dorey4		Crop: Winter Wheat				Year: 2020		
Field Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0,1,2,4,8)	Weight Factor	Weight Risk
Soil Erosion	NA	<5 tons/as/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	QA> 10 for erodible soils	0	1.5	0.0
Furrow Irrigation Erosion	N/A	Tail water recovery, QS>6 very erodible soils, or QS>10 other soils	QS> for erosion resistant soil	QS> for erodible soils	QA>6 for very erodible soils	0	1.5	0
Sprinkler Irrigation Erosion	All fields 0-3% slope, all sandy fields or field evaluation indicates little or no runoff large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8% large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	1.5	0.0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1 <input type="text"/>	0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	0	0.5	0.0
Commercial P Fertilizer Application Method	None Applied	Placed with Planter or injection deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporate d >3 months before crop or surface applied <3 months before crop emerges	Surface applied to pasture or >3 months before crop emerges	1 <input type="text"/>	1.0	1.0
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	1 <input type="text"/>	1.0	1.0

Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges	0	1.0	0.0
Organic P Source Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	0	1.0	0.0
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or application are directly into concentrated surface water flow areas.	0	1.0	0.0
<b>Total Phosphorus Index Value:</b>								<b>2.5</b>

**Appendix A: Phosphorus Index Worksheet (Complete for each field and crop)**

Field: E2								
Crop: Alfalfa Grass Mix								
Year: 2020								
Field Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0,1,2,4,8)	Weight Factor	Weight Risk
Soil Erosion	NA	<5 tons/as/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	QA> 10 for erodible soils	0	1.5	0.0
Furrow Irrigation Erosion	N/A	Tail water recovery, QS>6 very erodible soils, or QS>10 other soils	QS> for erosion resistant soil	QS> for erodible soils	QA>6 for very erodible soils	0	1.5	0
Sprinkler Irrigation Erosion	All fields 0-3% slope, all sandy fields or field evaluation indicates little or no runoff large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8% large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	1.5	0.0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1 <input type="text"/>	0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	0	0.5	0.0
Commercial P Fertilizer Application Method	None Applied	Placed with Planter or injection deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporate d >3 months before crop or surface applied <3 months before crop emerges	Surface applied to pasture or >3 months before crop emerges	8 <input type="text"/>	1.0	8.0
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	1 <input type="text"/>	1.0	1.0

Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges	0	1.0	0.0
Organic P Source Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	0	1.0	0.0
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or application are directly into concentrated surface water flow areas.	1	1.0	1.0
<b>Total Phosphorus Index Value:</b>								10.5

**Appendix A: Phosphorus Index Worksheet (Complete for each field and crop)**

Field: E3		Crop: Spring Wheat				Year: 2020		
Field Category Factor	None (0)	Low (1)	Medium (2)	High (4)	Very High (8)	Risk Value (0,1,2,4,8)	Weight Factor	Weight Risk
Soil Erosion	NA	<5 tons/as/yr	5-10 ton/ac/yr	10-15 tons/ac/yr	QA> 10 for erodible soils	0	1.5	0.0
Furrow Irrigation Erosion	N/A	Tail water recovery, QS>6 very erodible soils, or QS>10 other soils	QS> for erosion resistant soil	QS> for erodible soils	QA>6 for very erodible soils	0	1.5	0
Sprinkler Irrigation Erosion	All fields 0-3% slope, all sandy fields or field evaluation indicates little or no runoff large spray on silts 3-8%	Medium spray on silty soils 3-15% slopes, large spray on silty soils 8-15% slope, low spray on silt soils 3-8% large spray on clay soil 3-15% slope	Medium spray on clay soils 3-8% slopes, large spray on clay soils >15% slope, medium spray on silt soil >15% slope	Medium spray on clay soils >8% slope, low spray on clay soil 3-8% slope, low spray on silty soils >15% slopes	Low spray on clay soils >8% slopes	0	1.5	0.0
Runoff Class	Negligible	Very Low or Low	Medium	High	Very High	1 <input type="text"/>	0.5	0.5
Olson Soil Test P	-----	<20 ppm	20-40 ppm	40-80 ppm	>80 ppm	0	0.5	0.0
Commercial P Fertilizer Application Method	None Applied	Placed with Planter or injection deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporate d >3 months before crop or surface applied <3 months before crop emerges	Surface applied to pasture or >3 months before crop emerges	1 <input type="text"/>	1.0	1.0
Commercial P Fertilizer Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	2 <input type="text"/>	1.0	2.0

Organic P Source Application Method	None Applied	Injected deeper than 2 inches	Incorporated <3 months prior to planting or surface applied during growing season	Incorporated >3 months before crop or surface applied <3 months before crop.	Surface applied to pasture or >3 months before crop emerges	0	1.0	0.0
Organic P Source Application Rate	None Applied	<30 lbs/ac P205	31-90 lbs/ac P205	91-150 lbs/ac P205	>150 lbs/ac P205	0	1.0	0.0
Distance to Concentrated Surface Water Flow	>1,000 feet	200-1,000 feet, or functioning grass waterways in concentrated surface water	100-200 feet	<100 feet	0 feet or application are directly into concentrated surface water flow areas.	0	1.0	0.0
<b>Total Phosphorus Index Value:</b>								<b>3.5</b>